

Title: Flow-ecology relationships and environmental flows assessment for the Ozark-Ouachita Interior Highlands and the West Gulf Coastal Plains

Project Summary: Providing adequate water quantity and quality in streams and rivers is a pressing issue worldwide. Determining appropriate environmental flows in streams is critical for defining and designing landscapes capable of sustaining natural resources at desired levels. This proposal develops the second phase in a multi-year study, involving many partners and a series of steps towards the goal of producing the scientific basis for environmental flow standards within the Ozark-Ouachita Interior Highlands and the West Gulf Coastal Plains. Important products of this work will be regional flow-ecology relationships that will form the scientific framework for setting environmental flow standards and understanding impacts of land use and climate change. These flow-ecology relationships will help determine environmental flow needs in the Ozark-Ouachita Interior Highlands and the West Gulf Coastal Plains and will provide the basis for conservation of numerous aquatic species of greatest conservation need, including yellowcheek darter, Arkansas darter, least darter, Strawberry River darter, swamp darter, Ozark shiner, longnose darter, silver redhorse, paleback darter, goldstripe darter, current darter, Ozark chub, gapped ringed crayfish, coldwater crayfish, Neosho midget crayfish, Mammoth Spring crayfish, Meek's crayfish, ringed crayfish, William's crayfish, and numerous species of aquatic insects. This work will positively impact many species and ecosystems region-wide, those of greatest conservation need and otherwise.

Project Lead: Daniel D. Magoulick, Assistant Unit Leader and Professor, USGS, Arkansas Cooperative Fish & Wildlife Research Unit, Department of Biological Sciences, University of Arkansas, 479-575-5449; 479-575-3330Fax; danmag@uark.edu

Project Partners: Jeffrey W. Quinn, Stream Management Biologist (Statewide), Arkansas Game and Fish Commission, 213 A Highway 89 S, Mayflower, AR 72106, 877-470-3309, jwquinn@agfc.state.ar.us

Total SWG Funding Requested: \$94,201

Amount and Source of Matching Funds or In-kind Services: \$52,270

\$37,066 unrecovered indirect cost, University of Arkansas

\$15,204 graduate student tuition at University of Arkansas

Total Project Cost: \$146,471

Funding priority actions addressed:

- Determine environmental flow needs for aquatic communities
- Project will provide information for many SGCN species, including 12 fish, 7 crayfish and many insects (see Project Summary)

Ecobasins targeted: All basins within Ozark-Ouachita Interior Highlands and the West Gulf Coastal Plains (Fig. 1)

Need

Providing adequate water quantity and quality (i.e., environmental flow) in streams and rivers is vital for maintenance of adequate human and wildlife water supply and for maintaining function across ecosystems as lotic waters interface with terrestrial and marine ecosystems.

Environmental flows are viewed as a top environmental priority by many global organizations, yet determining adequate levels is often difficult because they can be impacted by many anthropogenic and natural factors, such as resource development (e.g. natural gas extraction), impoundment, irrigation and drought. Water use worldwide has increased steadily in the past 50 years (Holland 2007). Flows of water in streams and rivers affect freshwater organisms and ecosystems, but the relationship between them is often little studied and poorly known because it requires significant measurement of flow characteristics and biota at appropriate temporal and spatial scales. Knowledge of the flow magnitude, frequency, timing, duration, and rate of change needed to maintain particular organisms or ecosystem structure and function (environmental flows) would allow managers and conservation biologists to conserve high quality freshwater resources. Additionally, climate models predict temperatures and extreme precipitation events will increase in the Ozark-Ouachita Interior Highlands and West Gulf Coastal Plain regions (Diffenbaugh et al. 2005), so determining environmental flows would allow researchers and managers to assess potential impacts of climate change on stream organisms or ecosystem structure and function (Xenopoulos et al. 2005).

Several recent approaches for environmental flow determination exist. One of these approaches, Ecological Limits of Hydrologic Alteration (ELOHA) appears to have promise in determining flows needed to maintain particular organisms or ecosystem structure and function (Poff et al. 2010). Key aspects of the approach include using existing hydrological and biological databases, stakeholder involvement in setting goals and risk tolerance, and an adaptive management process. The large volume of region-wide biological data available from numerous sources provides a data-rich starting point to initiate ELOHA for the Ozark-Ouachita Interior Highlands and West Gulf Coastal Plain regions.

Additionally, the principal investigator recently completed a SWG-funded study examining flow-ecology relationships in a single flow regime within the Ozark Highlands (Leasure et al. 2014). Key findings of this study were: 1) Flow-ecology relationships varied temporally, 2) Water quality, geomorphology and habitat were often as, or more, important than hydrology, 3) Substantial variation among taxonomic groups in flow alteration-ecology relationships, and 4) Relatively high levels of uncertainty in flow-ecology relationships (even for significant relationships). Important remaining questions that need to be addressed are whether flow-ecology relationships vary spatially or by flow class? It is also important to improve our understanding of temporal variation in flow-ecology relationships given the substantial variation in flow-ecology relationships that we observed in just two years, as well as augmenting data sources to reduce our level of uncertainty.

The ultimate goal of our proposed research is to develop region-wide environmental flow standards. This proposal develops the second phase in a multi-year study, involving many partners and a series of steps towards the goal of producing the scientific basis for environmental flow standards within the Ozark-Ouachita Interior Highlands and West Gulf Coastal Plain regions. These steps include identifying and collating region-wide hydrological and biological databases, a hydrologic classification of rivers, and conducting aquatic community sampling at key sites in order to develop flow-ecology relationships within the Ozark-Ouachita Interior Highlands and West Gulf Coastal Plain regions. A hydrologic classification of the region has been completed as part of a SWG-funded study (Leasure et al. 2016), but additional work is needed to complete the rest of the process.

The relationships developed in the following objective can form the basis for setting regional environmental flow standards and understanding impacts of land use and climate change. Specifically, we will establish flow-ecology relationships for the Ozark-Ouachita Interior Highlands and West Gulf Coastal Plain regions. This will positively impact many species and ecosystems region-wide, those of greatest conservation need and otherwise. It will also be useful for addressing water use issues, such as those stemming from natural gas development.

Objective

Develop flow-ecology relationships for fish and macroinvertebrates for the Ozark-Ouachita Interior Highlands and West Gulf Coastal Plain regions using data from existing sources (e.g. USGS, EPA)

Approach

Database analysis to develop fish and macroinvertebrate flow-ecology relationships

A recently completed companion SWG-funded study had three main objectives: 1) To classify streams into natural flow regimes, 2) To calculate a hydrologic disturbance index at specific gauged and ungauged sites (Leasure et al. 2016, Fig. 1), and 3) To determine flow-ecology relationships in a single flow regime within the Ozark Highlands (Leasure et al. 2014).

Additionally, as part of a current SWG-funded study we have estimated flow alteration at gauged sites and we are developing models to predict flow alteration at ungauged sites within the Ozark-Ouachita Interior Highlands and West Gulf Coastal Plain regions. The present proposal intends to build on this work and will further develop flow-ecology relationships. First, we propose to identify and further collate hydrological (mainly USGS) and biological (e.g. USGS, EPA, USFS, TNC, state agencies) databases to develop a spatial relation between existing flow and biological inventory data within each flow regime in the Ozark-Ouachita Interior Highlands and West Gulf Coastal Plains. We hypothesize that flow alteration and hydrologic disturbance affects fish and macroinvertebrate population and community structure and dynamics. We will examine relationships between ecological response variables (e.g. species richness, diversity, density, traits) and flow alteration, hydrologic disturbance, geomorphic, land use, and climate variables to address this overarching hypothesis.

Hydrology data will be collected from the USGS National Water Inventory System. Ecologically relevant hydrological variables will be estimated from these data (Leasure et al. 2016, Fig. 2). Fish and macroinvertebrate biological databases for the Ozark-Ouachita Interior

Highlands and West Gulf Coastal Plains will be collected from all potential sources for initial screening. For both databases, an initial set of sites will be screened and a subset of sites will be selected based on data quality, temporal sequence, timing, drainage area, missing data, and other measures of data acceptability. We have already collected and collated some of this data, but we will add more data to this database for both hydrology and biology. We will examine all natural flow regimes within the Ozark-Ouachita Interior Highlands and West Gulf Coastal Plains (Leasure et al. 2016) with adequate hydrology and biology data for further analysis.

Limitations of using existing hydrology and biology databases to examine flow-ecology relationships are spatial and temporal gaps in data, poor data quality and lack of standardization, and mismatches between hydrological and biological data. Further, studies of this type are typically restricted to gauged sites, constraining the potential sample size. We will use flow alteration and hydrologic disturbance data estimated at un-gauged stream sites based on our current and previous research. This will allow us to greatly expand our use of existing biological data and alleviate two of the problems mentioned above. Additionally, we will use a new cutting edge approach, stream spatial network analysis, for dealing with data that shows lack of independence (i.e. most existing data). The principal investigator has attended a workshop on this approach and will train the graduate students to use this approach.

Expected Results and Benefits

Results from this work will provide the scientific foundation for ultimately producing environmental flow standards within the Ozark-Ouachita Interior Highlands and West Gulf Coastal Plains. We will identify and collate region-wide hydrological and biological databases and develop flow-ecology relationships for multiple flow regimes within the Ozark-Ouachita Interior Highlands and West Gulf Coastal Plains. The relationships developed can then form the basis for setting regional environmental flow standards and understanding impacts of land use and climate change. This work will positively impact many species and ecosystems region-wide (see Project Summary for list of SGCN affected). It will provide essential information and tools for dealing with water use issues, such as those resulting from natural gas development.

Principal investigator and students will share information from this project via publications and presentations at local to international meetings. As part of this project we will add data to the databases maintained by the Arkansas Game and Fish Commission and the Arkansas Wildlife Action Plan and the National Monitoring Partnership database. Finally, we will provide annual progress reports and a final report after the completion of the project.

Budget

Total SWG Funding Requested: \$94,201

Total Project Match: \$52,270

Total Project Cost: \$146,471

Requested SWG Funds	2016	2017	Total
Salary/Benefits			
2 Graduate Students	\$38,601	\$38,601	\$77,201
Operating Expenses			
Travel	\$4,000	\$4,000	\$8,000
Materials & Supplies	\$1,000		\$1,000

Capital Expenses	\$8,000		\$8,000
Total	\$51,601	\$42,601	\$94,201

Literature Cited

- Diffenbaugh, N. S., J. S. Pal, R. J. Trapp, and F. Giorgi. 2005. Fine-scale processes regulate the response of extreme events to global climate change. *Proceedings of the National Academy of Sciences of the United States of America* 102:15774-15778.
- Falcone, J.A., D.M. Carlisle, L.C. Weber. 2010. Quantifying human disturbance in watersheds: variable selection and performance of a GIS-based disturbance index for predicting the biological condition of perennial streams. *Ecological Indicators* 10:264-273.
- Holland, T.W. 2007. Water use in Arkansas, 2005. U.S. Geological Survey Scientific Investigations Report 2007–5241, 32 p.
- Leasure, D.R., D.T. Lynch and J.R. Schluterman. 2014. Classification of Arkansas flow regimes, regional flow-ecology relationships and environmental flows assessment for the Ozark region. Final Report prepared for the State Wildlife Grant program of the Arkansas Game and Fish Commission, Little Rock, Arkansas.
- Leasure, D.R., D.D. Magoulick and S. D. Longing. 2016. Natural flow regimes of the Ozark-Ouachita Interior Highlands region. *River Research and Applications* 32:18-35.
- Poff, N. L., B. D. Richter, A. H. Arthington, S. E. Bunn, R. J. Naiman, E. Kendy, M. Acreman, C. Apse, B. P. Bledsoe, M. C. Freeman, J. Henriksen, R. B. Jacobson, J. G. Kennen, D. M. Merritt, J. H. O'Keefe, J. D. Olden, K. Rogers, R. E. Tharme, and A. Warner. 2010. The ecological limits of hydrologic alteration (ELOHA): a new framework for developing regional environmental flow standards. *Freshwater Biology* 55:147-170.
- Xenopoulos, M. A., D. M. Lodge, J. Alcamo, M. Marker, K. Schulze, and D. P. Van Vuuren. 2005. Scenarios of freshwater fish extinctions from climate change and water withdrawal. *Global Change Biology* 11:1557-1564.

Project Leader

Dr. Daniel D. Magoulick - Assistant Unit Leader/ Professor, USGS, Arkansas Cooperative Fish and Wildlife Research Unit, Department of Biological Sciences, University of Arkansas 2000-present
 Ph.D. in Biological Sciences from University of Pittsburgh – 1994
 Author of 38 peer-reviewed scientific publications

Project Partners

Jeffrey W. Quinn, Stream Management Biologist, Arkansas Game and Fish Commission
 M.S. in Biology from University of Arkansas – 1998
 Author of 6 peer-reviewed scientific publications

Previous SWG Grants

Magoulick, D.D., R.J. DiStefano, B.K. Wagner and J.W. Fetzner, Jr. 2014-2016. Arkansas Game and Fish Commission. Invasive species effects, population status and population genetics

of crayfish species of greatest conservation need (*Orconectes marchandi*, *Orconectes eupunctus*, and *Cambarus hubbsi*) in the Ozark Highlands of Arkansas and Missouri. \$80,208.

Magoulick, D.D. and D.R. Leasure. 2014-2016. Arkansas Game and Fish Commission. Quantification of hydrologic alteration and relationships to biota in Arkansas streams: Development of tools and approaches for un-gaged streams. \$53,000.

Magoulick, D.D., S. Longing, J.W. Quinn, J. Jackson, J. Duzan and J. Petersen. 2011-2014. Arkansas Game and Fish Commission. Classification of Arkansas flow regimes, regional ecological-flow response relationships and environmental flows assessment for the Ozark region. \$172,000.

Magoulick, D.D. 2005-2007. Arkansas Game and Fish Commission. Effect of the introduced crayfish, *Orconectes neglectus*, on native crayfish in the Spring River drainage. \$68,390.

Figures




Figure 1. Natural flow regimes of 64 reference gages were identified using mixture- model cluster analysis based on 10 flow metrics. Natural flow regimes of all stream segments were predicted based on climate and catchment characteristics using a random forest model.




Figure 2. Flow metrics used in cluster analysis compared among natural flow regimes: Groundwater Stable (GS), Groundwater (G), Groundwater Flashy (GF), Perennial Runoff (PR), Runoff Flashy (RF), Intermittent Runoff (IR), and Intermittent Flashy (IF).




Figure 3. Important fish ecology-environment relationships in 2012. RHA is a qualitative index of stream geomorphology and habitat quality. Hydrologic disturbance index (HDI) is based on factors such as dam density, proximity to roads and canals, and water withdrawals (Falcone et al. 2010). Only significant relationships in the best models are shown. Blue lines indicate 95% confidence intervals.